



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Whence,

$$u^2 = \frac{k^2}{2} \left(\frac{1 - \sqrt{1 - c^2}}{\sqrt{1 - c^2}} \right),$$

and

$$v^2 = c^2(k^2 + u^2) = \frac{c^2 k^2}{2} \left(\frac{1 + \sqrt{1 - c^2}}{\sqrt{1 - c^2}} \right).$$

Hence,

$$u = \frac{k}{2\sqrt[4]{1 - c^2}} (\sqrt{1 + c} - \sqrt{1 - c}), \quad v = \frac{ck}{2\sqrt[4]{1 - c^2}} (\sqrt{1 + c} + \sqrt{1 - c}),$$

and

$$u(v - u) = \frac{k^2}{2} (1 - \sqrt{1 - c^2}).$$

The available work is h ft. lbs. per sec. per lb. of water used; the work utilized is

$$\frac{u}{g} (v - u) = h(1 - \sqrt{1 - c^2});$$

the proportion used is $(1 - \sqrt{1 - c^2})$.

If we let $c = \sin \alpha$, we have $u^2 = gh \tan \alpha \tan \alpha/2$, $v = 2u \cos^2 \alpha/2$, and efficiency = vers α .

331. Proposed by CLIFFORD N. MILLS, Brookings, S. Dakota.

A cyclist is riding due west at a speed of 12 miles per hr., and the wind is at the same time blowing from the southeast with a speed of $5\frac{1}{2}$ miles per hour. If the cyclist carries a small flag, in what direction will this flag fly? At what speed would the cyclist need to ride if the flag is to fly due north?

SOLUTION BY B. J. BROWN, Victor, Colo.

The component of the wind north is $11\sqrt{2}/4$ miles per hr., while that west is $11\sqrt{2}/4$ miles per hr. The resistance offered to the machine traveling west is $[12 - (11\sqrt{2}/4)]$ and the reaction is toward the east. Hence the forces affecting the flag are $[12 - (11\sqrt{2}/4)]$ toward east, and $11\sqrt{2}/4$ toward north and the direction θ , which the resultant makes with the east-west line = $\arctan 11\sqrt{2}/(48 - 11\sqrt{2}) = 25^\circ 36' 55''$ N. of E. In order for the flag to fly due N. the rider must travel W. at rate of $11\sqrt{2}/4$ miles per hr. Then he counteracts the resistance E. and only the component north is effective.

Also solved by PAUL CAPRON, W. J. THOME, and G. W. HARTWELL.

NUMBER THEORY.

236. Proposed by V. M. SPUNAR, Chicago, Illinois.

Find integral values of x, y, z , such that

$$xy + z = \square, \quad yz + x = \square, \quad zx + y = \square.$$

NOTE BY G. H. LING, University of Saskatchewan.

The following proposition generalizes somewhat the solution of this problem which was given in the February, 1917, MONTHLY.

THEOREM. If (1) a is any integer, (2) N_1, N_2 are integers such that $N_1 \cdot N_2 = a^2 + 1$, (3) $x = N_1 - 1, y = N_2 - 1, z = N_1 + N_2 - 1 + 2a$, then

$$(I) \quad xy + z = (a + 1)^2; \quad yz + x = (N_2 + a - 1)^2; \quad zx + y = (N_1 + a - 1)^2,$$

$$(II) \quad xy + x + y = a^2; \quad yz + y + z = (N_2 + a)^2; \quad zx + x + z = (N_1 + a)^2.$$

The solution given in the MONTHLY is the special case of this one in which

$$a = n^2 + n + 1.$$